arXiv:1210.0158v1 [nucl-th] 29 Sep 2012

# Probing in-medium spin-orbit potential with intermediate-energy heavy-ion collisions

Jun Xu[1, *] and Bao-An Li[1, 2, †]

[1]*Department of Physics and Astronomy, Texas A&M University-Commerce, Commerce, TX 75429-3011, USA*
[2]*Department of Applied Physics, Xi'an Jiao Tong University, Xi'an 710049, China*

Incorporating for the first time both the spin and isospin degrees of freedom explicitly in transport model simulations of heavy-ion collisions at intermediate energies, we observe that a strong local spin polarization appears in the early stage of the reaction. Most interestingly, it is found that the nucleon spin up-down differential transverse flow is a sensitive probe of the spin-orbit potential independent of its tensor terms, providing a novel approach to probe both the density and isospin dependence of the in-medium spin-orbit coupling that is important for understanding the structure of rare isotopes and synthesis of superheavy elements.

PACS numbers: 25.70.-z, 24.10.Lx, 13.88.+e

The spin-orbit coupling is common to the motion of many objects in nature from quarks, nucleons, and electrons to planets and stars. In nuclear physics, it has been very well known for a long time that the spin-orbit coupling is crucial for understanding the structure of finite nuclei, such as the magic numbers [1, 2]. However, many interesting questions regarding the in-medium spin-orbit coupling, especially its density and isospin dependence, remain unresolved although in free space it has been well determined from nucleon-nucleon (NN) scattering data [3]. Indeed, several phenomena observed or predicted in studying properties of nuclear structures have been providing us some useful information about the in-medium spin-orbit interaction. For instance, the kink of the charge radii of lead isotopes can only be explained by introducing a weakly isospin-dependent spin-orbit coupling [4, 5]. Moreover, strong experimental evidences of a decreasing spin-orbit coupling strength with increasing neutron excess were reported [6, 7]. Furthermore, new experiments are currently being carried out at several laboratories to explore the density and isospin dependence of the spin-orbit coupling by comparing energy splittings of certain orbits in the so-called "bubble" nuclei [8, 9] with those in normal nuclei [10]. The knowledge of the in-medium spin-orbit interaction is useful for understanding properties of drip-line nuclei [11], the astrophysical r-process [12], and the location of stability island for superheavy elements [13, 14]. On the other hand, it is also well known that the tensor force may affect the single-particle energy or even modify the magic numbers [15], and may change the evolution of energy gaps between certain orbits with isospin asymmetry of finite nuclei [16]. Moreover, calculations going beyond mean-field approximations indicate that the tensor force is also important for understanding short-range NN correlations [17, 18] and the density dependence of nuclear symmetry energy [19, 20]. Within Hartree-Fock calculations, the spin-orbit interaction leads to a density gradient term in the spin-orbit potential, while both the finite-range and tensor interactions contribute to the spin-orbit potential through the spin-current density terms, i.e., the tensor terms [21–23]. This mixing makes it very challenging to disentangle respective contributions from the spin-orbit interaction and the tensor force to phenomena found in nuclear structure studies. It will be extremely useful if one could find ways to study them separately and determine their respective strengthes. In this Letter, within a newly developed spin-isospin dependent transport model, we show that the nucleon spin up-down differential transverse flow in heavy-ion collisions at intermediate energies is a sensitive probe of the in-medium spin-orbit potential independent of its tensor terms.

Considering only time-even components of nuclear potentials, the single-particle Hamiltonian in nuclear matter can be written as [24]

$$h_q = -\frac{1}{2m}\nabla^2 + U_q + \vec{W}_q \cdot (-i\nabla \times \vec{\sigma}). \; (q = n, p) \quad (1)$$

In the above, the second and third terms are the central and spin-orbit potential, respectively. The time-odd components are neglected as including them would generally enhance dissipations [25, 26] but would not change the conclusion of the present work. Taking into account the density [27] and isospin dependence [4] as well as the tensor contribution [21], the form factor of the spin-orbit potential can be generally written as

$$\vec{W}_q = \frac{W_0^\star(\rho)}{2}(a\nabla\rho_q + b\nabla\rho_{q'}) + \alpha\vec{J}_q + \beta\vec{J}_{q'}, \; (q \neq q') \quad (2)$$

where $\rho_{n(p)}$ is the neutron (proton) density. $W_0^\star(\rho) = W_0(\rho/\rho_0)^\gamma$ represents the density-dependence of the spin-orbit coupling with $\rho = \rho_n + \rho_p$ and $W_0$ being the strength at saturation density $\rho_0$. Different values of $\gamma$, $a$, and $b$ can be used to mimic its density and isospin dependence. We notice that with $\gamma = 0$, $a = 2$ and $b = 1$ Eq. (2) reduces to the spin-orbit potential of the standard Skyrme-Hartree-Fock model. While equal values for $a$ and $b$, and a nonzero value for $\gamma$ were predicted for the effective spin-orbit potential within a relativistic mean-field model [5]. The spin-current density $\vec{J}$ is the antisymmetric vector part of the spin-current tensor $J_{\mu\nu}$ [25]. Neglecting the density dependence of $W_0^\star$, the spin-orbit coupling constant $W_0$ ranges from about 100 MeVfm$^5$ to 200 MeVfm$^5$ considering the uncertainty of the tensor terms [21], while the values of $\alpha$ and $\beta$ representing the strength of the tensor terms are still quite uncertain.

It is interesting to note that effects of the spin-orbit coupling were found earlier clearly in heavy-ion collisions at both low energies and relativistic energies. For instance, within the time-dependent Hartree-Fock (TDHF) calculations the nucleon spin-orbit potential was found to affect the fusion threshold energy [28] and leads to a local spin polarization [26, 29]. In non-central relativistic heavy-ion collisions at 200 GeV/nucleon the partonic spin-orbit coupling was found to lead to a global quark spin polarization [30]. As the beam energy increases from low to intermediate energies around a few hundred MeV/nucleon, while effects of the spin-orbit coupling are expected to grow due to the higher angular momentum available, effects of NN scatterings which are mostly Pauli blocked at low energies also become increasingly important. Depending on how the spin of each particle changes during NN scatterings, the spin polarization might be strongly affected by the latter in heavy-ion collisions at intermediate energies.

To model the spin-isospin dynamics in heavy-ion collisions at intermediate energies, we incorporate explicitly the spin degree of freedom and the spin-orbit potential in a previously developed isospin-dependent Boltzmann-Uehling-Uhlenbeck (BUU) transport model, see, e.g., Refs. [31, 32]. The new model is dubbed SI-BUU12. To our best knowledge, the spin degree of freedom was never considered before in any of the existing transport models for heavy-ion reactions at intermediate energies since the emphasis of the community has been on extracting information about the Equation of State (EOS) of symmetric nuclear matter, density dependence of nuclear symmetry energy and in-medium NN scattering cross sections using spin-averaged experimental observables. In the SI-BUU12 model, the nucleon density $\rho$, spin density $\vec{s}$, and the spin-current density $\vec{J}$ are all calculated by using the test particle method [33, 34]. The equations of motion with the spin-orbit potential are

$$\frac{d\vec{r}}{dt} = \frac{\vec{p}}{m} + \vec{\sigma} \times \vec{W}_q, \tag{3}$$

$$\frac{d\vec{p}}{dt} = -\nabla U_q - \nabla \left[ \vec{W}_q \cdot (\vec{p} \times \vec{\sigma}) \right], \tag{4}$$

$$\frac{d\vec{\sigma}}{dt} = 2(\vec{W}_q \times \vec{p}) \times \vec{\sigma}. \tag{5}$$

In the center-of-mass (c.m.s) frame, two Fermi spheres of projectile/target nucleons are Lorentz boosted in the $\pm z$ direction. During the collision process, the form factor $\vec{W}$ points mainly along the impact parameter of the reaction, i.e., the x axis. Due to the spin-orbit potential, the nucleon spin $\vec{\sigma}$ tends to be parallel to $\vec{p} \times \vec{W}$ in order to lower the energy of the system. Namely, the nucleon spin is more likely to be polarized along the y axis perpendicular to the reaction plane (x-o-z). We will refer in the following text a nucleon with its spin in the +y (-y) direction as a spin-up (spin-down) nucleon. While our qualitative conclusions in this work are insensitive to the specific choice for $U_q$, numerical results presented in the following are obtained by using a momentum-independent potential corresponding to a soft EOS of incompressibility $K_0 = 230$ MeV, a symmetry energy $E_{sym} = 30$ MeV and its density slope $L = 60$ MeV at saturation density $\rho_0 = 0.16$ fm$^{-3}$ similar to the MSL0 parameterization [35].

Nucleon-nucleus scattering experiments have shown that nucleons may flip their spins after NN scatterings due to spin-related nuclear interactions, see, e.g., Ref. [36] for a review. Although not well determined yet, the spin-flip probability for in-medium NN scatterings is known to be appreciable, depending on the collision energy and the momentum transfer [37]. In the present work we will test different options of setting spins, including randomizing, flipping, or keeping spins unchanged after each NN scattering to study effects of different spin-flip probabilities on final observables. In addition, a spin- and isospin-dependent Pauli blocking is introduced in the SI-BUU12 model.



FIG. 1: (color online) Contours of nucleon reduced density $\rho/\rho_0$ (first row), y component of spin density $s_y$ (second row), and x component of the density gradient $(\nabla\rho)_x$ (third row) and spin-current density $J_x$ (fourth row) in the reaction plane at different stages in Au+Au collisions at a beam energy of 400 MeV/nucleon and an impact parameter of b = 8 fm. Note that the x coordinates of the projectile (target) is negative (positive).

As an illustration of spin-orbit coupling effects, it is interesting to first examine for a typical reaction at intermediate energies the time evolution of the y component of the spin density $s_y$, the x component of the density gradient $(\nabla\rho)_x$ and the spin-current density $J_x$ in comparison with the nucleon density contour in the reaction plane. Shown in Fig. 1 are these quantities for Au+Au collisions at a beam energy of 400 MeV/nucleon and an impact parameter of b = 8 fm. For this example, we set $W_0 = 130$ MeVfm$^5$, $\gamma = 0$, $a = 2$, $b = 1$, $\alpha = \beta = 0$, and the spins of the colliding nucleons are randomized after each scattering. We will only change one or two of the above parameters while keeping all the rest the same in the following. Initially, we put the projectile and target nuclei without spin polarization far away so that a stable local spin polarization due to the spin-orbit potential can be generated in the two nuclei before they collide in the c.m.s frame. The initial distance is set to be long enough so that its further increment will not affect much the spin polarization of the two nuclei when they get in touch around 60 fm/c. Indeed, a local spin polarization is induced in the early stage of the reaction as expected. This "spin twist" phenomenon was first observed in TDHF calculations [26, 29]. We note that the induced spin polarization observed here is stronger than that at lower energies due to the higher nucleon momentum at intermediate energies. It is seen that participant nucleons are more likely to spin-down while the spectator nucleons are more likely to spin-up during the early stage of the reaction. In the later stage, however, the spin polarization becomes weaker because of NN scatterings and other spin mixing effects, especially for participant nucleons at higher densities. At the end of the reaction, except for some large spectator-like fragments, the local spin polarization vanishes for most nucleons.

As shown in the equations of motion, the spin-orbit coupling also affects the nucleon momentum and spatial distributions. In particular, spin-up and spin-down nucleons with the same momentum experienced opposite spin-orbit potentials during the whole collision process. Nucleon transverse collective flow, measured by using the average transverse momentum $< p_x(y) >$ in the reaction plane versus rapidity $y$, is one of the best known observable for revealing effects of density gradients in nuclear reactions [38]. What can we learn about the spin-orbit potential from the nucleon transverse flow? To answer this question, first we compare in the left panel of Fig. 2 the transverse flows of spin-up and spin-down nucleons. It is seen that the transverse flow of spin-up nucleons is larger than that of spin-down ones. This can be easily understood by looking at the x-component of the density gradient shown in the third row of Fig. 1. By examining the time evolution, we found that the effects of the spin-orbit potential on the transverse flow during the first 20 fm/c of the collision are mostly washed out due to violent interactions. The spin-dependent transverse flow is mainly determined by the dynamics afterwards. As the projectile (target) is still moving in the +z (-z) direction, the participant nucleons from the projectile (target) with negative (positive) $(\nabla\rho)_x$ give a more repulsive/attractive spin-orbit potential $[\vec{W} \cdot (\vec{p} \times \vec{\sigma}) > / < 0]$ for spin-up/down nucleons. This leads to a larger transverse flow for spin-up nucleons than spin-down ones.



FIG. 2: (color online) Transverse flow of spin-up nucleons and spin-down nucleons (a) and spin up-down differential flows using different $W_0$ (b) for the same reaction as in Fig.1.

To extract more accurately information about the spin-orbit potential without much hinderance of central potentials, we investigate next the spin up-down differential transverse flow

$$F_{ud}(y) = \frac{1}{N(y)} \sum_{i=1}^{N(y)} \sigma_i (p_x)_i, \tag{6}$$

where $N(y)$ is the number of nucleons with rapidity $y$, and $\sigma_i$ is $1(-1)$ for spin-up (spin-down) nucleons. Similar to the neutron-proton differential transverse flow for probing the symmetry potential [39], the spin up-down differential transverse flow maximizes effects of the opposite spin-orbit potentials for spin-up and spin-down nucleons while canceling out largely spin-independent contributions. Indeed, the spin up-down transverse flow is a sensitive probe of the spin-orbit coupling strength $W_0$. As an example, shown in the right panel of Fig. 2 is a comparison of the spin up-down differential transverse flows obtained using the two limiting values of $W_0$ used in the literature. As expected, a larger value of $W_0$ gives a stronger spin-orbit potential and thus a larger spin up-down differential transverse flow.

The density dependence of the spin-orbit coupling, which was tested earlier, see, e.g., Ref. [27], is still almost completely unknown, and this has recently motivated more new experiments [10]. To investigate effects of the density dependence of the spin-orbit coupling on the spin up-down differential flow, shown in the panel (a) of Fig. 3 are the results obtained by varying only the $\gamma$ parameter. It is seen that the spin up-down differential flow is larger for a weaker density dependence of the spin-orbit coupling if the strength at saturation density is fixed.

The isospin dependence of the spin-orbit coupling is another interesting issue especially relevant for under-



FIG. 3: (color online) Spin up-down differential transverse flows using different density dependence of the spin-orbit coupling coefficients (a) and from pure like-nucleon (b) and pure unlike-nucleon coupling (c) for the same reaction as in Fig.1.



FIG. 4: (color online) Comparison of the spin up-down differential transverse flows using different coefficients of the tensor terms (a) and different treatments of spin after scatterings (b).

standing the structure of rare isotopes and the synthesis of superheavy elements. We notice that available studies on nuclear structure seem to favor a weak isospin dependence of the spin-orbit coupling [4, 5, 11]. To evaluate potential applications of our approach in further constraining the isospin dependence of the spin-orbit potential, we next compare the spin up-down differential flows for neutrons and protons using the pure like-nucleon coupling ($a = 3$ and $b = 0$) and pure unlike-nucleon coupling ($a = 0$ and $b = 3$) in panel (b) and panel (c) of Fig. 3, respectively. As the system considered is globally neutron-rich and $\nabla\rho_n$ is generally larger than $\nabla\rho_p$, the pure like(unlike)-nucleon coupling leads to an appreciably larger (smaller) spin up-down differential flow for neutrons than for protons. Moreover, the unlike-nucleon coupling generally reduces significantly the overall strength of the spin-orbit potential and thus the spin up-down differential flow. Of course, more neutron-rich systems will be better for probing the isospin-dependence of the spin-orbit coupling using the double differential flow between spin up-down neutrons and protons.

So far we haven't included in the spin-orbit potential contributions from the tensor terms yet. With spin polarization, as $\vec{\sigma}$ is parallel to $\vec{p}\times\nabla\rho$, $\vec{J} \sim \vec{p}\times\vec{\sigma}$ is generally opposite to $\nabla\rho$. Thus, $\vec{J}$ terms with positive (negative) coefficients contribute negatively (positively) to $\vec{W}$, as mentioned in Ref. [29]. Although great efforts have been made in determining the values of $\alpha$ and $\beta$ from studying the structure of finite nuclei [21–23], they are still quite uncertain. By setting $\alpha = \beta = 100$ or $-100$ MeVfm$^5$, which are considered reasonable in magnitude based on the early studies in the literature, we compare the spin up-down differential transverse flow in the left panel of Fig. 4. It is seen clearly that the spin up-down differential transverse flow is rather insensitive to the $\vec{J}$ terms. This is understandable as the $\nabla\rho$ terms related to the density distribution remain strong while the spin-current density vanishes after the first 20 fm/c of the collision as shown in Fig. 1.

Finally, what are the effects of the possible spin flip in NN scatterings on the spin up-down differential flow? We answer this question by using the results shown in the right panel of Fig. 4. Due to the lack of knowledge about the energy and isospin dependence of the in-medium spin-flip probability, we compare the following three choices for setting spins of nucleons after NN scatterings: (1) flipped, (2) randomized, or (3) unchanged, effectively from a large to small spin-flip probability. It is seen that the spin up-down differential transverse flow decreases with the increasing spin-flip probability as one expects. Moreover, it is very encouraging to see that the spin up-down differential flow is still considerable even if a 100% spin-flip probability is assumed.

In summary, the spin degree of freedom and the spin-orbit potential are incorporated for the first time in an isospin-dependent transport model providing a useful new tool for investigating the spin-isospin dynamics of heavy-ion collisions at intermediate energies, such as the development of local spin polarization in these reactions. The nucleon spin up-down differential transverse flow is shown to be a sensitive probe of the in-medium spin-orbit potential independent of its tensor terms. Comparisons with future experiments will allow us to determine the density and isospin dependence of the in-medium spin-orbit coupling that has significant ramifications in both nuclear physics and astrophysics.

We thank Dr. Y. Iwata for useful communications and Dr. W. G. Newton and Dr. F. J. Fattoyev for helpful discussions. This work was supported in part by the US National Science Foundation grants PHY-0757839 and PHY-1068022 as well as the National Aeronautics and Space Administration under grant NNX11AC41G issued through the Science Mission Directorate.

---

[*] Electronic address: Jun.Xu@tamuc.edu
[†] Electronic address: Bao-An.Li@tamuc.edu

[1] M. Goeppert-Mayer, Phys. Rev. **75** 1969 (1949).
[2] O. Haxel, *et al.*, Phys. Rev. **75**, 1766 (1949).
[3] R. B. Wiringa, *et al.*, Phys. Rev. C **51**, 38 (1995).
[4] M. M. Sharma, *et al.*, Phys. Rev. Lett. **74**, 3744 (1995).
[5] P. G. Reinhard and H. Flocard, Nucl. Phys. **A584**, 467 (1995).
[6] J. P. Schiffer, *et al.*, Phys. Rev. Lett. **92**, 162501 (2004).
[7] L. Gaudefroy, *et al.*, Phys. Rev. Lett. **97**, 092501 (2006).
[8] B. G. Todd-Rutel, *et al.*, Phys. Rev. C **69**, 021301 (R) (2004).
[9] M. Grasso, *et al.*, Phys. Rev. C **79**, 034318 (2009).
[10] O. Sorlin, `http://www.uni-mainz.de/Organisationen/vistars/talks_russbach2012/russb12_sorlin.pdf`
[11] G. A. Lalazissis, *et al.*, Phys. Lett. **B418**, 7 (1998).
[12] B. Chen, *et al.*, Phys. Lett. **B355**, 37 (1995).
[13] M. Bender, *et al.*, Phys. Rev. C **60**, 034304 (1999).
[14] M. Morjean, *et al.*, Phys. Rev. Lett. **101**, 072701 (2008).
[15] T. Otsuka, *et al.*, Phys. Rev. Lett. **95**, 232502 (2005).
[16] T. Otsuka, *et al.*, Phys. Rev. Lett. **97**, 162501 (2006).
[17] R. Subedi, *et al.*, Science **320**, 1476 (2008).
[18] L. Frankfurt, *et al.*, Int. J. Mod. Phys. A **23**, 2991 (2008).
[19] S. O. Bäckman, *et al.*, Phys. Rep. **124**, 1 (1985).
[20] C. Xu and B. A. Li, Phys. Rev. C **81**, 064612 (2010).
[21] T. Lesinski, *et al.*, Phys. Rev. C **76**, 014312 (2007).
[22] G. Colò, *et al.*, Phys. Lett. **B646** 227 (2007).
[23] M. Bender, *et al.*, Phys. Rev. C **80**, 064302 (2009).
[24] D. Vautherin and D. M. Brink, Phys. Rev. C **5**, 626 (1972).
[25] Y. M. Engel, *et al.*, Nucl. Phys. **249**, 215 (1975).
[26] J. A. Maruhn, *et al.*, Phys. Rev. C **74**, 027601 (2006).
[27] J. M. Pearson and M. Farine, Phys. Rev. C **50**, 185 (1994).
[28] A. S. Umar, *et al.*, Phys. Rev. Lett. **56**, 2793 (1986).
[29] Y. Iwata and J. A. Maruhn, Phys. Rev. C **84**, 014616 (2011).
[30] Z. T. Liang and X. N. Wang, Phys. Rev. Lett. **94**, 102301 (2005); Phys. Lett. **B629**, 20 (2005).
[31] B. A. Li, *et al.*, Phys. Rev. Lett. **78**, 1644 (1997).
[32] B. A. Li, *et al.*, Phys. Rep. **464**, 113 (2008).
[33] C. Y. Wong, Phys. Rev. C **25**, 1460 (1982).
[34] G. F. Bertsch and S. Das Gupta, Phys. Rep. **160**, 189 (1988).
[35] L. W. Chen, *et al.*, Phys. Rev. C **82**, 024321 (2010).
[36] G. G. Ohlsen, Rep. Prog. Phys. **35**, 717 (1972).
[37] W. G. Love and M. A. Franey, Phys. Rev. C **24**, 1073 (1981); **27**, 438 (1983).
[38] P. Danielewicz and G. Odyniec, Phys. Lett. **B157**, 146 (1985).
[39] B. A. Li, Phys. Rev. Lett. **85**, 004221 (2000).